

A Geometrical Description of the “Classical” Weak-Field Hall Effect in Metals

F. D. M. Haldane

Department of Physics, Princeton University, Princeton NJ 08544-0708

(Dated: April 12, 2005)

Within the “anisotropic relaxation time” approach to transport processes, I obtain a new geometric formula for the weak-field Hall conductivity of metals, which generalizes previously-known formulas due to Tsuji (restricted to cubic 3D metals) and Ong (for 2D metals).

PACS numbers: 72.15.Lh, 72.15.Eb

While the Kubo formula in principle provides a fully-Hamiltonian description of the conductivity of metals, practical calculations generally use the simpler linearized semiclassical Boltzmann equation description, commonly combined with the additional “relaxation time” approximation[1]. This leads to a “geometric” expression for the DC Ohmic conductivities, which can be expressed in terms of the Fermi-surface geometry and a temperature-dependent “scattering path length” (mean free path) $\ell(\mathbf{s}, T) = v_F(\mathbf{s})\tau(\mathbf{s}, T)$ where $\mathbf{s} = \{s^1, s^2\}$ is a pair of curvilinear coordinates parameterizing points on the Fermi surface $\mathbf{k}_F(\mathbf{s})$, $v_F(\mathbf{s})$ is the Fermi speed, and $\tau(\mathbf{s}, T)$ is a notional temperature-dependent “anisotropic” ($\mathbf{s}$ -dependent) relaxation time for the quasiparticle distribution to relax back into equilibrium once the current-driving field is removed.

The phenomenological “scattering path length” can also be expressed as a vector mean free path, $\boldsymbol{\ell}(\mathbf{s}) = \ell(\mathbf{s})\hat{\mathbf{n}}_F(\mathbf{s})$, where $\hat{\mathbf{n}}_F(\mathbf{s})$ is the outward normal of the Fermi surface (direction of the Fermi velocity). The detailed microscopic origin of $\ell(\mathbf{s}, T)$ is usually rather inaccessible: theoretically, within the Boltzmann formalism, one must distinguish between “quasiparticle lifetimes”, and “transport lifetimes”, but the uncertainties of the microscopic relaxation mechanisms turn the scattering path length into a phenomenological fitting parameter. (See Ref.[2] for a review of the application of Boltzmann theory to metals.) It is probably most meaningful in the case where the dominant relaxation process is “catastrophic” inelastic scattering, so that all memory of the previous quasiparticle state is erased by the scattering process.

In this formalism, with unbroken time-reversal symmetry, the symmetric DC Ohmic conductivity tensor is

$$\sigma_{\Omega}^{ab}(T) = \frac{e^2}{\hbar} \sum_{\alpha} g_s \int_{S_{\alpha}} \frac{d^2 S_F}{(2\pi)^3} \hat{n}_F^a(\mathbf{s}) \hat{n}_F^b(\mathbf{s}) \ell(\mathbf{s}, T), \quad (1)$$

where $g_s = 2$ is the spin degeneracy factor, S_{α} are the topologically-distinct sheets (compact 2-manifolds) of the Fermi surface (labeled by α), and $d^2 S_F$ is the (scalar) Fermi-surface area element. To make the structure of geometrical quantities clearer, I use a “covariant” notation where spatial coordinates r^a , $a = 1, 2, 3$, have “upper” indices, and the dual k -space coordinates k_a have

“lower” indices, with the summation convention on repeated upper-lower pairs; this shows up the formal difference between $\mathbf{k} \cdot \mathbf{r} = k_a r^a$ which is metric-independent, and $\hat{\mathbf{n}}_F \cdot \hat{\mathbf{n}}_F = g_{ab}^0 \hat{n}_F^a \hat{n}_F^b = 1$, where $g_{ab}^0 = “\delta_{ab}” = \text{diag}(1, 1, 1)$ is the Euclidean metric tensor with $\det g^0 = 1$. In such a notation, the components of the electric field E_a have lower indices, while components B^a of the magnetic flux density have upper indices. In a metal with cubic symmetry, the conductivity tensor is isotropic, $\sigma_{\Omega}^{ab}(T) \propto g_0^{ab}$, where g_0^{ab} is the inverse of the Euclidean metric tensor:

$$\sigma_{\Omega}^{ab} = \frac{1}{3} g_0^{ab} \frac{e^2}{\hbar} \sum_{\alpha} g_s \int_{S_{\alpha}} \frac{d^2 S_F}{(2\pi)^3} \ell(\mathbf{s}, T), \quad (\text{cubic}). \quad (2)$$

If time-reversal symmetry is broken by the application of a weak uniform magnetic flux density $\mathbf{B}$, σ^{ab} acquires an antisymmetric Hall term σ_H^{ab} : to linear order in $\mathbf{B}$,

$$\sigma^{ab}(T, \mathbf{B}) = \sigma_{\Omega}^{ab}(T) + \frac{e^2}{\hbar} \epsilon^{abc} \gamma_{cd}(T) \left(\frac{eB^d}{\hbar} \right), \quad (3)$$

where ϵ^{abc} is the usual antisymmetric rank-3 Levi-Civita symbol, and $\gamma_{ab}(T)$ is a symmetric tensor. Using the same Jones-Zener solution[1] of the Boltzmann equation in the relaxation-time approximation that is usually used to derive (1), Tsuji[3] found a geometric expression for the weak-field Hall conductivity of cubic metals, which is equivalent to

$$\gamma_{ab} = \frac{1}{3} g_{ab}^0 \sum_{\alpha} g_s \int_{S_{\alpha}} \frac{d^2 S_F}{(2\pi)^3} H(\mathbf{s}) \ell^2(\mathbf{s}, T), \quad (\text{cubic}), \quad (4)$$

where $H(\mathbf{s})$ is the *mean curvature* of the Fermi surface at point $\mathbf{s}$. The Tsuji formula has been quantitatively tested[4] against band-structure calculations of the Fermi surface curvatures of various cubic metals, plus “reasonable” assumptions about $\tau(\mathbf{s}, T)$ (and hence $\ell(\mathbf{s}, T)$); that study indeed found sensitivity to Fermi surface regions of high mean curvature.

Unlike (1), the formulas (2) and (4) are not truly “local” Fermi surface formulas, because they essentially perform an integral of the average over the “star” of 48 equivalent points of a general Fermi surface point in a system with cubic symmetry. The purpose of this Letter is to give the full non-cubic generalization of Tsuji’s

formula as a true integral over local Fermi-surface properties; this new “classical” Hall effect formula complements another new geometric Fermi-surface formula[5] for the intrinsic “anomalous” ($\mathbf{B} = 0$) Hall conductivity of ferromagnetic metals, which is a fundamental quantum effect associated with quasiparticle Berry phases.

The Fermi-surface curvature field $\kappa^{ab}(\mathbf{s})$ is a symmetric tensor relating two “one-forms”, the differential change $d\hat{\mathbf{n}}_F$ in outward normal vector and the differential change $d\mathbf{k}_F$ of the Fermi vector for small displacements $d\mathbf{s}$ on the Fermi surface:

$$d\hat{\mathbf{n}}_F^a = \kappa^{ab}(\mathbf{s}) dk_{Fb}. \quad (5)$$

In a Euclidean coordinate frame that locally diagonalizes it at Fermi surface point $\mathbf{s}$, $\kappa^{ab}(\mathbf{s}) = \text{diag}(k_1^{-1}, k_2^{-1}, 0)$, and $n_F^a(\mathbf{s})n_F^b(\mathbf{s}) = \text{diag}(0, 0, 1)$, where k_1 and k_2 are the two principal Fermi-surface radii of curvature. Then

$$H(\mathbf{s}) = \frac{1}{2} \left(\frac{1}{k_1} + \frac{1}{k_2} \right), \quad K(\mathbf{s}) = \frac{1}{k_1 k_2}, \quad (6)$$

are respectively the mean (extrinsic) curvature and the Gaussian (intrinsic) curvature. Note that $(H(\mathbf{s}))^2 \geq K(\mathbf{s})$, so the mean curvature can only change sign (from electron-like $H(\mathbf{s}) > 0$ to hole-like $H(\mathbf{s}) < 0$) in regions of negative Gaussian curvature. The relation (5) can be also be expressed as

$$dk_{Fa} = k_{ab}(\mathbf{s}) d\hat{\mathbf{n}}_F^b, \quad (7)$$

where $k_{ab}(\mathbf{s}) = \text{diag}(k_1, k_2, 0)$ is the *radius-of-curvature field*, which is a well-defined symmetric tensor wherever the Gaussian curvature $K(\mathbf{s})$ is non-zero. It diverges when $K(\mathbf{s}) \rightarrow 0$, but the product $K(\mathbf{s})k_{ab}(\mathbf{s}) \equiv \epsilon_{ace}\epsilon_{bdf}\hat{n}_F^c(\mathbf{s})\hat{n}_F^d(\mathbf{s})\kappa^{ef}(\mathbf{s})$ remains finite.

The Gaussian curvature $K(\mathbf{s})$ has a profound geometric significance; it relates two fundamental “two-forms”, the Fermi surface (scalar) area element $d^2 S_F = g_{ab}^0 \hat{n}_F^a \epsilon^{bcd} dk_{Fc} \wedge dk_{Fd}$, and $d^2 \Omega_F = \epsilon_{abc} \hat{n}_F^a d\hat{\mathbf{n}}_F^b \wedge d\hat{\mathbf{n}}_F^c$, the solid angle subtended by the normal vector $\hat{\mathbf{n}}_F(\mathbf{s})$ as it varies over that area element:

$$K(\mathbf{s}) d^2 S_F = d^2 \Omega_F. \quad (8)$$

This leads to the Gauss-Bonnet theorem that

$$\frac{1}{4\pi} \int_{S_\alpha} K(\mathbf{s}) d^2 S_F = \frac{1}{4\pi} \int_{S_\alpha} d^2 \Omega_F = 1 - g_\alpha, \quad (9)$$

where the non-negative integer g_α is the *genus* (number of “handles”) of a Fermi-surface manifold S_α .

The genus is the primary topological characteristic of a Fermi surface manifold; another important topological property is the the dimension d_α^G of the Bravais Lattice of reciprocal vectors $\mathbf{G}$ generated by periodic open orbits on S_α . For a closed path Γ , $\oint_\Gamma d\mathbf{k}_F = \mathbf{G}(\Gamma)$, where $\mathbf{G}(\Gamma)$ is a non-zero reciprocal lattice vector if the image of Γ in

3D k -space is a periodic “open orbit”. On a 2-manifold of genus g_α , all closed paths Γ are topologically equivalent to paths built up as sequences of paths chosen from a set of $2g_\alpha$ elementary non-trivial closed paths Γ_i (the basis of generators of the homology group). The generators Γ_i can be chosen so that only $d_\alpha^G \leq 3$ of them have non-zero $\mathbf{G}(\Gamma_i)$, which are linearly independent, and form a basis of a d_α^G -dimensional Bravais lattice. A sphere-like “nearly-free-electron” 3D Fermi surface, as in the alkali metals, has $d_\alpha^G = 0$, a cylinder-like quasi-2D Fermi surface has $d_\alpha^G = 1$, a plane-like quasi-1D Fermi surface has $d_\alpha^G = 2$, and a multiply-connected Fermi surface with cubic symmetry, as in the noble metals, has $d_\alpha^G = 3$.

My main result, the non-cubic generalization of the Tsuji formula, can now be given:

$$\gamma_{ab}(T) = \sum_\alpha \frac{g_s}{2} \int_{S_\alpha} \frac{K(\mathbf{s}) d^2 S_F}{(2\pi)^3} k_{ab}(\mathbf{s}) \ell^2(\mathbf{s}, T). \quad (10)$$

Unlike (4), this is a true integral of local quantities on the Fermi surface, making no use of any point-group symmetries, and is expressed in a fully-covariant tensor formulation, independent of choice of coordinate frame. In the cubic case, the replacement $\gamma_{ab} \rightarrow \frac{1}{3} g_{ab}^0 g_0^{cd} \gamma_{cd}$ may be made, and the cubic Tsuji form (4) is recovered because $g_0^{ab} k_{ab}(\mathbf{s}) \equiv 2H(\mathbf{s})/K(\mathbf{s})$.

The derivation of (10) is as follows; let $n_0(\mathbf{k})$, for $\mathbf{k}$ near the Fermi surface in 3D k -space, be the ground-state Fermi-Dirac distribution of quasiparticle occupations in an independent-quasiparticle Ansatz, and let $\delta n_E(\mathbf{k}) = (e\tau(\mathbf{k})/\hbar) \mathbf{E} \cdot \nabla_k n_0(\mathbf{k})$ be the first-order non-equilibrium change induced by the uniform applied electric field $\mathbf{E}$, where ∇_k is the k -space derivative, and $\tau(\mathbf{k})$ is an appropriate phenomenological temperature-dependent anisotropic relaxation time. Here $\nabla_k n_0(\mathbf{k}) = \hbar \mathbf{v}_F(\mathbf{k}) f'(\delta\varepsilon(\mathbf{k}))$, where $f(\varepsilon)$ is the Fermi-Dirac distribution function $(\exp(\varepsilon/k_B T) + 1)^{-1}$, $\delta\varepsilon(\mathbf{k})$ is the quasiparticle excitation energy, and $\mathbf{v}_F(\mathbf{k}) = \hbar^{-1} \nabla_k \delta\varepsilon(\mathbf{k})$ is the quasiparticle group velocity. Then

$$\delta n_E(\mathbf{k}) = f'(\delta\varepsilon(\mathbf{k})) e \mathbf{E} \cdot \boldsymbol{\ell}, \quad (11)$$

where $\mathbf{v}_F(\mathbf{k})\tau(\mathbf{k}) \equiv \boldsymbol{\ell}(\mathbf{k}) \equiv \boldsymbol{\ell}(\mathbf{k})\hat{\mathbf{n}}_F$. Because it induces k -space motion parallel to the Fermi surface, a weak uniform magnetic flux density only produces a second-order change $\delta n_{E,B}(\mathbf{k}) = (e\mathbf{v}_F(\mathbf{k})\tau(\mathbf{k})/\hbar) \times \mathbf{B} \cdot \nabla_k \delta n_E(\mathbf{k})$:

$$\delta n_{E,B}(\mathbf{k}) = f'(\delta\varepsilon(\mathbf{k})) \boldsymbol{\ell} \times (e\mathbf{B}/\hbar) \cdot \nabla_k (e\mathbf{E} \cdot \boldsymbol{\ell}). \quad (12)$$

The current density is $g_s V^{-1} \sum_k e \mathbf{v}_F(\mathbf{k}) \delta n(\mathbf{k})$, where V is the volume of the system; in the Fermi-liquid regime, $f'(\delta\varepsilon(\mathbf{k}))$ vanishes except near the Fermi surface, and the expression for the induced current becomes a Fermi surface integral. The first-order term δn_E gives the Ohmic conductivity $\sigma_\Omega^{ab}(1)$; $\delta n_{E,B}$ gives the Hall term σ_H^{ab} as

$$\frac{e^3}{\hbar^2} \sum_\alpha g_s \int_{S_\alpha} \frac{d^2 S_F}{(2\pi)^3} (\hat{n}_F^a \boldsymbol{\ell}) (\mathbf{B} \times \hat{\mathbf{n}}_F \cdot \nabla_k) (\hat{n}_F^b \boldsymbol{\ell}).$$

This is an antisymmetric tensor in indices a, b , as

$$d^2 S_F(\mathbf{B} \times \hat{\mathbf{n}}_F \cdot \nabla_k) \equiv B dk_{F\parallel} dk_{F\perp} \partial / \partial k_{F\perp},$$

where $dk_{F\parallel}$ and $dk_{F\perp}$ are components of $d\mathbf{k}_F$ parallel and perpendicular to $\mathbf{B}$; the symmetrized form of the expression for σ_H^{ab} is the integral of the derivative of a periodic function over a full period, and thus vanishes. After explicitly antisymmetrizing the expression, one can replace $\nabla_k(\hat{\mathbf{n}}_F^b \ell)$ by $\ell(\nabla_k \hat{\mathbf{n}}_F^b)$, and then $\mathbf{B} \times \hat{\mathbf{n}}_F \cdot \nabla_k \hat{\mathbf{n}}_F^b$ by $\epsilon_{cde} B^c \hat{\mathbf{n}}_F^d \kappa^{eb}$; finally the identity $\epsilon_{ace} \epsilon_{bdf} \hat{\mathbf{n}}_F^c \hat{\mathbf{n}}_F^d \kappa^{ef} \equiv K k_{ab}$ is used to obtain (10).

There have been previous attempts to generalize the Tsuji formula (e.g., Ref.[6]); these did not uncover a general geometric formula with the simplicity of (10) because they typically expressed the solution of the Boltzmann equation in terms of non-geometric quantities such as higher k -space derivatives of the quasiparticle dispersion $\delta\varepsilon(\mathbf{k})$. What is remarkable about (10) (and (1)) is that they only involve Fermi surface geometry and the scattering path length, with no direct reference to Fermi-liquid energy parameters such as $v_F(\mathbf{s})$. As in general relativity, the use of a covariant metric-independent and coordinate-frame-independent tensor formulation greatly aids the identification of formulas with the correct geometric structure.

There is a subtle change of the interpretation of the Fermi surface in the formulas presented here. Instead of thinking of the Fermi surface as a set of 2D surfaces embedded in a periodically-repeated 3D k -space, it is best to view it as a set of compact 2-manifolds $\mathcal{S}_\alpha$ on which quasiparticle states are defined, and which are the domains of functions such as $\mathbf{k}_F(\mathbf{s})$, $\hat{\mathbf{n}}_F(\mathbf{s})$, $k_{ab}(\mathbf{s})$, etc. On such a 2-manifold, any “two-form” can be used to formulate a surface integral. In particular, $\ell^2 K d^2 S_F = \ell^2 d^2 \Omega_F = \ell^{-1} \ell \cdot d^2 \mathbf{S}_\ell$, where $(d^2 \mathbf{S}_\ell)_a$ is the (oriented) vector surface area element $\epsilon_{abc} d\ell^b \wedge d\ell^c$ of the (temperature-dependent) 2D surface swept out by $\ell(\mathbf{s}, T)$ as $\mathbf{s}$ varies over $\mathcal{S}_\alpha$.

This leads to an alternative form of (10), inspired by the novel approach of Ong[7], who studied the Hall conductivity σ_H^{2D} of a 2D metal where quasiparticle motion takes place only on electronically-isolated lattice planes with no interplane tunneling, and where the Fermi surface is a set of 1-manifolds Γ_α , so $\mathbf{k}_F(\mathbf{s})$ is a one-parameter curve in the planar Brillouin zone. Ong pointed out that, instead of focusing on $\mathbf{k}_F(\mathbf{s})$, it was far more informative to examine the planar curve traced out by the 2D mean-free-path vector $\ell_0(\mathbf{s}, T)$, and obtained

$$\sigma_H^{2D}(T) = \frac{e^2}{h} \sum_\alpha \frac{g_s}{2} \oint_{\Gamma_\alpha} \frac{d\ell_0}{(2\pi)^2} \cdot \ell_0 \times \frac{e\mathbf{B}}{h}, \quad (13)$$

which expresses the 2D Hall conductivity of a single plane *completely* in terms of $\ell_0(\mathbf{s}, T)$. In analogy to Ong’s approach, the 3D formula (10) can also be written as

$$\gamma_{ab}(T) = \sum_\alpha \frac{g_s}{2} \int_{\mathcal{S}_\alpha} \frac{\ell \cdot d^2 \mathbf{S}_\ell}{(2\pi)^3} k_{ab}(\mathbf{s}) \ell^{-1}(\mathbf{s}). \quad (14)$$




FIG. 1: (Left) 3D k -space image (Fermi vector $\mathbf{k}_F(\mathbf{s})$), and (right) the corresponding 3D ℓ -space image (vector mean-free-path $\ell(\mathbf{s})$) of a quasi-2D Fermi surface (quasiparticle 2-manifold parametrized by $\mathbf{s}$). $\mathbf{G}^0$ is the primitive reciprocal vector that indexes the weakly-coupled lattice planes. $\ell(\mathbf{s})$ is parallel to the outward Fermi-surface normal $\hat{\mathbf{n}}_F(\mathbf{s})$. Both images are of a surface with genus $g = 1$, an intrinsic property independent of the coordinates ($\mathbf{k}_F(\mathbf{s})$ or $\ell(\mathbf{s})$) used to represent the 2-manifold. In the 2D limit, the toroid $\ell(\mathbf{s})$ collapses into a planar loop $\ell_0(\mathbf{s})$ with $\mathbf{G}^0 \cdot \ell_0 = 0$.

If a 3D system is formed by a family of isolated metallic 2D lattice planes indexed by a primitive reciprocal lattice vector $\mathbf{G}^0$, $\sigma_H^{ab} = \sigma_H^{2D} \epsilon^{abc} (G_c^0 / 2\pi)$, with $\mathbf{G}^0 \cdot \ell_0 = 0$. Then the 2D formula (13) is equivalent to

$$\gamma_{ab}(T) = G_a^0 \epsilon_{bcd} \sum_\alpha \frac{g_s}{2} \oint_{\Gamma_\alpha} \frac{d\ell_0^c}{(2\pi)^3} \ell_0^d, \quad (2D \text{ limit}). \quad (15)$$

Then (15) can be obtained from (14) in the limit that a quasi-2D Fermi surface with genus $g = 1$ (and $d^G = 1$) degenerates into a strictly-2D system (see Fig.1). The limiting process is not entirely straightforward, because in the 2D limit, the Gaussian curvature $K(\mathbf{s}) \rightarrow 0$ everywhere on the Fermi surface. The k -space image of such a Fermi surface is a weakly-modulated cylindrical surface connecting two opposite Brillouin-zone faces (relatively-displaced by $\mathbf{G}^0$), while the ℓ -space image is a toroid, with one small radius that shrinks to zero in the 2D limit, so the toroid collapses into a loop $\ell_0(\mathbf{s})$ in the plane $\ell_0 \cdot \mathbf{G}^0 = 0$.

For passage to the 2D limit, the vector mean-free-path $\ell(\mathbf{s})$ can be written as $\ell_0(\mathbf{s}) + \delta\ell(\mathbf{s}, s_\perp)$, where $\ell_0 \cdot \mathbf{G}^0 = 0$, and $\delta\ell \cdot \mathbf{G}^0$ depends only on $s_\perp$; here, both $\mathbf{s}$ and $s_\perp$ are periodic parametric Fermi surface coordinates, so displacements $d\mathbf{k}_F$ with constant $s_\perp$ are normal to $\mathbf{G}^0$, and those with constant $\mathbf{s}$ are along the component of $\mathbf{G}^0$ parallel to the Fermi surface. In the 2D limit, $\delta\ell(\mathbf{s}) \rightarrow 0$, but $k_{ab}(\mathbf{s})$ has a divergent part $\Lambda(\mathbf{s}) G_a^0 G_b^0$, where $\Lambda \rightarrow \infty$. Any path along which $s_\perp$ increases at fixed $\mathbf{s}$ is a periodic open orbit with $\oint ds_\perp k_{ab} \partial n_F^b / \partial s_\perp = \mathbf{G}_a^0$; so as $\delta\ell \rightarrow 0$, $\oint ds_\perp \Lambda(\partial / \partial s_\perp) \delta\ell \cdot \mathbf{G}^0 \rightarrow \ell_0(\mathbf{s})$, which remains finite. In the 2D limit, $\ell \cdot d^2 \mathbf{S}_\ell \rightarrow (\ell_0 \times d\ell_0) \cdot (\partial \delta\ell / \partial s_\perp ds_\perp)$. Since $\ell_0 \times d\ell_0$ is parallel to $\mathbf{G}^0$, (14) is now easily found to evolve into (15).

In 2D, σ_H^{2D} has dimensions [resistance] $^{-1}$, and is $e^2/\hbar$ times a dimensionless factor that is invariant under rescaling of the Fermi surface. In 3D, in contrast, σ_H^{ab} has dimensions [resistance] $^{-1} \cdot$ [wavenumber] and scales linearly with Fermi-surface scale; thus the 3D formula (14) depends on Fermi-surface geometry through the radius-of-curvature field $k_{ab}(\mathbf{s})$, and not just on the vector mean-free-path $\ell(\mathbf{s}, T)$ alone, unlike the 2D formula (13).

There is an important difference between the images of the Fermi surface 2-manifold as represented by the Fermi vector field $\mathbf{k}_F(\mathbf{s})$ and the vector mean-free-path field $\ell(\mathbf{s})$. In k -space, “level repulsion” means that the Fermi surface $\mathbf{k}_F(\mathbf{s})$ is generically non-self-intersecting (except in exceptional non-symmorphic space-group cases where it crosses certain Brillouin zone boundaries). Even point degeneracies, associated with intersections with axes of k -space rotation symmetry (or rare stable “accidental” degeneracies), are not common. In contrast, ℓ -space surfaces $\ell(\mathbf{s})$ have no self-avoidance property, and may well intersect in complicated ways.

The formulas (10) and (14) presented here are “weak field” (small- $\mathbf{B}$) formulas. The “weak-field” condition is usually characterized as “ $\omega_c \tau \ll 1$ ”. The appropriate geometrical condition is that the Fermi-surface displacement $\Delta \mathbf{k}_F = \ell(\mathbf{s}) \times (e\mathbf{B}/\hbar)$ as the quasiparticle travels its mean free path is such that the change $\Delta \hat{\mathbf{n}}_F$ in its real-space direction-of-motion is small, where

$$(\hat{\mathbf{n}}_F \times \Delta \hat{\mathbf{n}}_F)_a = \ell(\mathbf{s}, T) K(\mathbf{s}) k_{ab}(\mathbf{s}) \left(\frac{eB^b}{\hbar} \right). \quad (16)$$

From a geometric Fermi-liquid point of view, the scattering path length $\ell(\mathbf{s}, T)$ may be a “more fundamental” quantity than the quasiparticle lifetime $\tau(\mathbf{s}, T)$. It controls the maximum possible real-space extent of a Gaussian quasiparticle wavepacket, as coherence would be lost if the size was larger. By the uncertainty principle, $\ell(\mathbf{s}, T)^{-1}$ is a lower bound on the quasiparticle localization near the Fermi surface in k -space. Assuming that “natural” Gaussian quasiparticle wavepackets should have quasi-isotropic spreads in k -space, this local k -space lengthscale may also provide a natural “quantum geometric” scale for non-orthogonality of wavepackets centered at neighboring Fermi surface points, and may play the role of a (temperature-dependent) “quantum length” analogous to the magnetic length $\ell_B = \sqrt{\hbar/|eB|}$ in lowest-Landau level physics.

It seems appropriate to characterize a transport regime in which the new weak-field Hall conductivity formula

las (10), (14) are valid as a “classical” Hall regime, as it seems to involve no aspect of the quasiparticle dynamics other than the underlying geometrical constraint provided by the Fermi surface, which defines a (3+2)-dimensional symplectic phase-space structure that, at each $\mathbf{s}$, singles out one real-space direction ($\hat{\mathbf{n}}_F(\mathbf{s})$) as “special” (not conjugate to a displacement $d\mathbf{k}_F$ on the Fermi surface). Equilibrium fluctuations driven by “catastrophic” inelastic scattering events nucleate a quasiparticle at Fermi surface point $\mathbf{s}$, and it propagates a distance $\ell(\mathbf{s}, T)$ in the local “special” direction before another violent inelastic event causes its decoherence, and returns the charge to the local “reservoir”. During its lifetime, the quasiparticle’s trajectory can be influenced by electric and magnetic fields. This seems to be an entirely “classical” diffusive process, with no involvement of quantum interference or Hamiltonian dynamics, as none of the microscopic ingredients of Fermi liquid theory (Fermi speed $v_F(\mathbf{s})$, Landau functions, quasiparticle Berry phases[5], *etc.*) are referenced in (10), (14).

In summary, I have described a geometric formulation of the “classical” weak-field Hall effect in metals (within a relaxation-time approach), and given two equivalent forms (10), (14) of a new formula expressing the Hall conductivity entirely in terms of Fermi surface geometry and a quasiparticle mean-free-path length that is traversed between scattering events that erase memory of its prior history. This generalizes previously-known formulas that were restricted to special cases: Tsuji’s formula[3] for metals with cubic symmetry, and Ong’s formula[7] for 2D metals.

This work was supported in part by the U. S. National Science Foundation (under MRSEC Grant No. DMR02-13706) at the Princeton Center for Complex Materials.

-
- [1] H. Jones and C. Zener, Proc. R. Soc. A **145**, 268 (1934).
 - [2] P. B. Allen, in *Quantum Theory of Real Materials*, edited by J. R. Chelikowsky and S. G. Louie (Kluwer, Boston, 1996), Chap. 17.
 - [3] M. Tsuji, J. Phys. Soc. Jpn. **13** 979 (1958).
 - [4] W. W. Schulz, P. B. Allen, and N. Trivedi, Phys. Rev. B **45**, 10886 (1992).
 - [5] F. D. M. Haldane, Phys. Rev. Lett. **93**, 206602 (2004).
 - [6] W. Kesternich, Phys. Rev. B **13**, 4227 (1976).
 - [7] N. P. Ong, Phys. Rev. B **43**, 193 (1991).